J. S. DEPARTMENT OF COMMERCE

R. P. LAMONT, Secretary

COAST AND GEODETIC SURVEY

R. S. PATTON, Director

Special Publication No. 159

THE BOWIE METHOD OF TRIANGULATION ADJUSTMENT

AS APPLIED TO THE FIRST-ORDER NET IN THE WESTERN PART OF THE UNITED STATES

BY

OSCAR S. ADAMS

SENIOR MATHEMATICIAN

275 .435 no.159



UNITED STATES
GOVERNMENT PRINTING OFFICE
WASHINGTON: 1930

National Oceanic and Atmospheric Administration

ERRATA NOTICE

One or more conditions of the original document may affect the quality of the image, such as:

Discolored pages
Faded or light ink
Binding intrudes into the text

This has been a co-operative project between the NOAA Central Library and the Climate Database Modernization Program, National Climate Data Center (NCDC). To view the original document, please contact the NOAA Central Library in Silver Spring, MD at (301) 713-2607 x124 or www.reference@nodc.noaa.gov.

LASON Imaging Contractor 12200 Kiln Court Beltsville, MD 20704-1387 January 1, 2006

PUBLICATION NOTICE

The Coast and Geodetic Survey maintains mailing lists containing the names and addresses of persons interested in its publications. When a new publication

and addresses of persons interested in its publications. When a new publication or a new edition is issued on any of the subjects covered by the mailing list, a circular showing its scope and contents is sent to each person whose name and address is on the mailing list of the subject covered by the publication.

If you desire to receive notices regarding publications of the Coast and Geodetic Survey as issued, write to the Director, Coast and Geodetic Survey, Washington, D. C., indicating the mailing lists on which you wish your name entered, or, if you prefer, check the list on the form below, remove this sheet and mail it to the Director, Coast and Geodetic Survey, Washington, D. C.

	D-4-
The DIRECTOR HAUTER STATES (Date Coast and Geodetic Survey,
THE DIRECTOR, UNITED STATES	Washington, D. C.
check below to receive notices a Geodetic Survey: [] 109. Astronomic Work [] 109-B. Coast Pilot. [] 109-B. Coast Pilot. [] 109-D. Geodesy, or Mea [] 109-E. Gravity. [] 109-F. Hydrography. [] 109-H. Nautical Charts. [] 109-H. Nautical Charts. [] 109-J. Precise Traverse. [] 109-L. Terrestrial Magn [] 109-M. Tides. [] 109-M. Topography. [] 109-O. Triangulation.	name be placed on the mailing lists indicated by regarding publications issued by the Coast and c. surement of the Earth.
[] 109-P. Cartography.	
	Name
	Address

CONTENTS

· · · · · · · · · · · · · · · · · · ·
Abstract
Introduction
Growth of triangulation net
Need for general adjustment
Bowie method devised
Description of the plan of adjustment
Description of the plan of adjustment
Typical junction figures
Preliminary adjustment of sections
Computation of preliminary positions
Adjustment of latitude and longitude closures
Formation of observation equations
Solution of normal equations
Computations
ComputationsAssumed positions of junction stations
Fixed nosition
Fixed positionComputed differences of position of junction stations
Observation equations
Table for formation of latitude normal equations
Table for formation of longitude normal equations.
Normal equations for latitude
Normal equations for langitude
Normal equations for longitude
Solution of normal equations Back solution for latitude
Back solution for latitude
Back solution for longitude
Latitude and longitude v's and total closures
Changes in positions of junction points caused by the readjustment
Closures from the adjustment.
Changes in positionsSection closures
Section closures
Kind of figures
Conclusion
ILLUSTRATIONS Fig.
1. Simple junction figure containing base and Laplace azimuth
2. Complicated junction figure involving several stations.
3. Junction figure which contains no bases or Laplace azimuths
4. Diagram of net showing numbers assigned to junction points
4. Diagram of net snowing numbers assigned to junction points
5. Loop closures resulting from readjustment
6. Section closures resulting from apportionment of loop closures
7. Changes of positions of junction points, previously adjusted, caused by
the readjustment

THE BOWIE METHOD OF TRIANGULATION ADJUSTMENT AS APPLIED TO THE FIRST-ORDER NET IN THE WESTERN PART OF THE UNITED STATES

By OSCAR S. ADAMS, Senior Mathematician, U. S. Coast and Geodetic Survey

ABSTRACT

This publication contains a discussion of the new method for the adjustment of a triangulation net as applied in the western part of the United States. extent of the arcs and their complicated interrelations made it practically impossible to adjust the whole with one set of equations; accordingly a new method of handling such networks was devised employing junction figures and sections of arcs between them. A complete discussion of the procedure in the application of the method in the western half of the United States is given, together with an account of the method of establishing the junction figures. The formation of the equations for the loop closures is given in detail and the method of least squares is employed to determine the best positions to be held in the junction squares is employed to determine the best positions to be held in the junction figures.

The tables give the results of the preliminary computations and the equations derived from the loop closures. The formation and solution of the normal equations are also given to serve as the examples for such applications. Lastly the final results derived by the solutions are given also as a table showing the changes in position of previously adjusted junction points.

At the end of the publication there is given a discussion of the facts disclosed

by this extensive adjustment of triangulation arcs. This section also includes a discussion regarding the accuracy of the observations as disclosed by the remarkable closures of loops throughout the whole area.

INTRODUCTION

The first-order triangulation of the United States has been extended from year to year with a more or less degree of regularity for the past 80 years or more. The arcs in the western part of the United States have been in process of extension for approximately 50 years. This more or less uniform growth consisted of extending new ares from those already in existence. As the various arcs were observed they were adjusted by least squares so that the work should be consistent when the results were furnished for engineering applications. When any loop was completed, the total discrepancy in closure was adjusted into the last arc that closed the loop. This quite frequently imposed unfavorable conditions upon the last arc and resulted in distortions of the work that were not justified by the general accord of the observations. For practical reasons this method had to be followed, for otherwise the whole system of triangulation would have to be readjusted every time a new loop was closed. The economic conditions thus necessitated that this plan should be followed until the main chains of triangles should be completed and then a unified adjustment would be undertaken for the whole region.

The final arcs for the framework in the western part of the United States were completed in 1926, and then the general adjustment was

carried through. The plan upon which the adjustment should be made was decided upon in 1924 and many of the preliminary computations were made before the final arcs were completed. An undertaking of such proportions could only be carried through by the aid of a large body of men trained in such calculations. Especial mention should be made of a number of members of the division of geodesy; W. F. Reynolds, C. H. Swick, H. C. Mitchell, W. D. Sutcliffe, and Howard S. Rappleye aided in the preparation of the equations for the adjustments, as well as by giving expert advice upon technical points. In the further calculations the main work was done by G. L. Fentress, H. P. Kaufman, F. W. Darling, J. A. Duerksen, W. Shofnos, C. J. Clifford, and W. E. Wood.

GROWTH OF TRIANGULATION NET

In the adjustment of triangulation it is evident that the closure of a loop should be distributed throughout its entire length. This is a very simple matter when the arc consists of a single loop such as that formed by carrying triangulation around an island or around a lake. However, when the system consists of a number of interconnected loops, the solution of the equations of condition for all of the loops in a single set presents serious difficulty on account of the amount of computation that would be required in carrying out the

adjustment.

In the United States the geodetic datum was adopted when the triangulation system consisted of a meager skeleton joining the arc near the Atlantic coast to that near the Pacific coast. As this mere outline was supplemented by new work, the discrepancies in the closures of loops were adjusted into the new arcs, the part already adjusted being held fixed. In the earlier work no true geodetic azimuths, derived by applying the Laplace correction to observed astronomic azimuths, were held in the adjustments. Experience had not yet shown that an arc of triangulation tends to swerve in azimuth unless it is held in orientation by equations of condition derived from the observed azimuths corrected for the deflection of the vertical. This departure from true orientation in the previously adjusted arcs at times caused comparatively large discrepancies to appear when new arcs were observed to close loops. As the country became divided up by many closed arcs of triangulation, this method of adjustment became more and more objectionable because often comparatively short arcs were forced to absorb loop closures that were out of all proportion to their lengths, and as a result the corrections that had to be applied to them were unduly large. In all loops the discrepancies disclosed were such as might be expected from the grade of the work when the whole length of the loop was considered, but they were unduly large for adjustment into the new work with old work held fixed. It became evident that at some time a readjustment would have to be made of the whole, or at least part, of the net to relieve this condition of unequal strain. Since the western part of the country was divided up into about the ideal-sized loops, the part from the ninety-eighth meridian arc inclusive was chosen for the first application of the method.

NEED FOR GENERAL ADJUSTMENT

Although this section was sufficiently covered by arcs of triangulation to permit of a general adjustment that would give consistent results which could then be held fixed for all time, yet a serious practical difficulty arose when the adjustment was contemplated. If the adjustment should be made by the method of least squares including the whole region in one adjustment, it would involve the solution of more than 3,000 simultaneous equations. Even with the use of modern calculating machines this would require an enormous amount of work, much more than could be justified economically. There would also be the possibility of some error in the formation or the solution that would fail to be detected until the final results were being computed, and then it would require much extra labor to make the corrections.

The question, however, was an important one at that time, for the arc of first-order triangulation, extending from the northwestern part of the United States along the coast of British Columbia and through southeastern Alaska, was awaiting adjustment, and this could not be undertaken until final values for the geographic positions of the stations from which the new arc would extend could be obtained. This arc was measured and adjusted by the cooperation of this bureau and the Geodetic Survey of Canada. The data carried by this arc into Alaska are the controlling data for the whole of the work in Alaska as well as for the work in western and northwestern Canada. It thus was imperative that the best value possible should be derived for the line in northwest Washington from which the Canadian-Alaskan arc extended.

BOWIE METHOD DEVISED

An economically feasible method for making an adjustment of such an extensive net of triangulation was worked out early in 1924 by Dr. William Bowie, chief of the division of geodesy of the United States Coast and Geodetic Survey. He conceived and formulated a plan by means of which such an extensive system can be adjusted with sufficient rigor at comparatively small cost and in a comparatively short time. By this Bowie method the discrepancies in loop closures of a net formed by arcs of triangulation are distributed in a way somewhat similar to that used in the adjustment of a first-order level net.

The triangulation in the western part of the United States consists of a number of intersecting arcs. In order to make the readjustment of the net by the new method, the unit was taken as the section of an arc between two junction points. Hereafter in this publication this

unit will be called simply a "section."

This same method of adjustment can also be applied to an area completely covered by triangulation by selecting loops formed by chains of well-shaped triangles, quadrilaterals, or other figures with the intermediate figures or stations omitted. Such omitted stations can later be adjusted into the general net by supplementary computations.

After the loop closures have been distributed among the various sections of the arcs of triangulation forming the loops, and in this way the most probable geographic positions of the stations used as junction points have been derived, then the sections can be adjusted

separately between the junction figures.

At the request of Doctor Bowie, the writer worked out a method for forming the equations and for making the adjustments by the method of least squares which are necessary to render the plan practicable. On first consideration it was contemplated that the adjustment might be made by the use of interrelated equations for the loop closures, but on further investigation it was found that no economical scheme of this kind could be devised. After careful study a plan was perfected for the use of two independent sets of equations, one set for the closures in latitude and the other set for the longitude closures. This plan was found to be both economical and practical and at the same time it gives a very satisfactory distribution of the

discrepancies in loop closures.

After working out the mathematical theory of the method, the author was put in entire charge of the execution of the project. From 8 to 12 mathematicians were employed on the adjustments of the sections, the number varying as the availability of the work demanded. As a result, in about 15 months after starting the computations the work had advanced to the point where the adjustment of the loop closures could be made to obtain the most probable positions of the junction points to be held in the junction figures. One of the great advantages of the Bowie method is the fact that a large number of computers can be employed upon the work at the same time. When the plan of the adjustment was initiated not all of the field work of the various arcs to be included was completed. all of the observational data had been ready, as many computers could have been put upon the computations as there were sections, and the final results could have been obtained in a much shorter time. As the matter stood with this bureau, efforts were made to complete the work as soon as all of the field work was available. The record for accomplishment speaks well both for the Bowie method and for the efficiency of the mathematicians who were employed upon the various computations necessary to carry out the scheme. We believe that this accomplishment deserves the attention of all practical geodesists, showing as it does a new method for bringing about an adjustment of complicated nets of triangulation in a comparatively short time with sufficient rigor to meet all demands, either practical or scientific. The economy both of time and expense is a fact to consider as well as the fact that the method is applicable to adjustments of triangulation that could scarcely be handled in any other way. In actual working days for one man the time employed up to the determination of the junction points was 2,200 days of 7 hours each. For the computations of the sections between junction figures the time consumed was 1,600 days, making a total of 3,800 days of 7 hours each for the whole work of the adjustment of the net.

DESCRIPTION OF THE PLAN OF ADJUSTMENT

In carrying out the scheme of adjustment the first step consisted in laying out junction figures at the various intersections of the arcs. The general aim was to make these figures as simple as possible but to choose them, if possible, so that the various sections radiating from them should be attached to the junction figure by a single line. It was not possible in all cases to attain this ideal, but it was so arranged whenever feasible. The junction figures in some cases consisted of only one or two quadrilaterals, but in other cases they

became fairly complicated figures. If the plan had been in mind during the time when the arcs were being observed, no doubt the ideal simple figures could have been observed in almost every case. However, since the work was already done it remained to make the best use of what was at hand.

After the junction figures were laid out, it then became necessary to adopt some length and azimuth to hold in them. The ideal case was a junction figure in which was included a measured base and an observed Laplace azimuth. The figure could then be adjusted by least squares, and the measured base and azimuth could furnish the required control. Each line from which a section started was fixed in length and azimuth by this adjustment and these values were then

used when the adjoined section was adjusted.

If either a measured length or an observed Laplace azimuth was wanting in the junction figure, it became necessary to adopt a value for some line of the figure for length or azimuth, or for both in case neither was present in the figure. This was generally done by setting up equations from the nearest control data through each of the radiating sections to some line in the junction figure and then some weighted mean was adopted as the value to be used. The weights were assigned by consideration of the strength of the various connections, a base or azimuth lying near the line having more weight than one farther away. The main object was to adopt control data that would best accord with the nearest bases and Laplace azimuths in the various sections. It was always possible to arrive at some reasonable compromise among the various values derived from the equations in the sections. In some cases where the chosen junction figure had already been adjusted in one of the arcs, the adjusted values of the angles and lengths were held fixed if they agreed reasonably well with the lengths derived from the other sections. Various methods were thus used in fixing these data, with the sole object in mind to adopt values that would best fit in with the bases and Laplace azimuths in the radiating sections.

LAPLACE AZIMUTHS

Since there may be some question as to how the Laplace azimuth values were obtained, it is probably best to give some account of them at this point. A Laplace azimuth is an observed astronomical azimuth that has been corrected for the effect of the deflection of the vertical. This correction can be computed if an astronomical longitude has been observed at the station as well as the astronomical azimuth. If we denote the astronomical value by A and the geodetic value by G, we have the relation,

$$(A-G)$$
 in azimuth = $-[(A-G)$ in longitude] $\sin \phi$

in which ϕ is the latitude of the station. In computing this correction the best available value for the geodetic longitude as derived from the preliminary computations is used. This will not, in most cases, differ from the final value by more than a few tenths of a second of arc, and hence the computed correction will be very nearly correct. After the final value of the geodetic longitude is adopted in the junction figure, a small correction to the azimuth—as held can be

made to take care of the change in the geodetic longitude, and this change in the azimuth can be taken care of in the final adjustment of the sections between the junction figures.

TYPICAL JUNCTION FIGURES

As examples of the various conditions that may arise in the work of establishing the fundamental length and azimuth in junction figures, three diagrams are included showing three typical cases that arose in this adjustment. Figure 1 shows an example of the simplest case in which a measured base (indicated by a heavy line) and a Laplace azimuth (indicated by an arrowhead) are included in a simple figure. In Figure 2 we also have these fundamental data included in the junction figure, but the figure itself is considerably more complicated. Finally, in Figure 3 we have an example of the most complicated case. The length and azimuth had to be carried into the junction figure from four radiating sections. The simple chain of triangles through which the data had to be carried is shown in each case. The junction figure itself was somewhat complicated even




FIGURE 1.—Simple junction figure containing base and Laplace azimuth

after these accessory triangles were omitted. This was the most involved case that arose in the whole adjustment. In any case of the application of this method of adjustment the length and azimuth in the junction figure could be determined in any reasonable way that might be desired; the end in view should be to choose these data so as best to accord with the other fixed data in the adjoining sections. In the present adjustment this aim was kept in view at all times and the actual method of attaining it was varied at times to suit the individual case.

PRELIMINARY ADJUSTMENT OF SECTIONS

· After a junction figure has been adjusted, the lengths and azimuths The sections can of the lines to which the sections join are known. then be adjusted between the junction figures holding these lengths and azimuths. All of the conditions will thus be included except the conditions for closure in latitude and longitude. As a matter of expediency the latitude and longitude equations were made up in this preliminary adjustment with no closure term added to them. "forward" solution of the normal equations could thus be made with these equations carried along as the last two equations of the set. In the preliminary "back" solution these two equations were ignored. However, after the discrepancy in closure was later determined, it was only necessary to carry the forward solution through these equations, neglecting the constant terms of the other normals except for a small correction to the constant term of the last azimuth equation of each section due to the change in the Lapace azimuth caused by the change in the longitude. In this way an additional set of corrections were derived which, added algebraically to the preliminary corrections, gave the final complete corrections. In this way an

economy in the labor required was attained. However, in the application of the Bowie method, after the final positions in the junction figures have been determined (in the manner to be indicated later), a complete new adjustment of each section, including all of the conditions at one time, could be made if so desired. It would require considerable more work, however, and the results would not be any more satisfactory. Economy of calculation should have some weight in deciding which course to follow. There is no question but that the method used in the present adjustment saved a great deal of time and consequent expense.

The preliminary adjustment is no more than the usual figure adjustment including the angle equations, the side equations, the length equations to hold the lengths of the measured bases and of the fixed lines at the ends of the section, and the azimuth equations to hold any intermediate Laplace azimuths and the azimuths of the junction All possible conditions are thus included except the conditions of closure in

latitude and longitude.

COMPUTATION OF PRELIMINARY **POSITIONS**

After these preliminary adjustments have all been made the geographic positions can be computed, starting from the point whose latitude and longitude is to be held fixed as the fundamental datum. These preliminary positions can be computed along any of the arcs and, consequently, there will be certain junction stations at which there will be discrepancies of closure. If computations are to be carried further from such a junction one or the other of the consistent set of data Thus the positions of all should be used. stations can be traced in unbroken line through some continuous arcs back to the fundamental station.

From these positions an assumed position is taken for some chosen point in each of




FIGURE 2.-GURE 2.—Complicated junction figure involving several stations

the junction figures. The formation of the observation equations is simplified if these assumed positions are made to correspond in each case with the preliminary positions as computed through one of the continuous arcs. In this way the constant term in many of the equations will be zero and the work of formation of the latitude and longitude equations will be simplified thereby.

ADJUSTMENT OF LATITUDE AND LONGITUDE CLOSURES

After the positions have been assumed for the junction points, the observation equations are formed for both the latitude closures and the longitude closures. The procedure is the same in each case.

The difference between the assumed latitudes of the junction points at the two ends of a section gives the assumed value for this difference. By subtracting from this quantity the computed value for the same we get an equation for the v in latitude for that section; a similar procedure gives the v in longitude for the section. The constant term in these will be expressed in units of seconds of arc. Seconds of arc, however, represent different linear lengths depending upon the lati-





FIGURE 3.—Junction figure which contains no bases or Laplace azimuths

The lengths in the junction figure at the intersection of the arcs are controlled by the four bases indicated by the heavy lines and the azimuths by the four Laplace azimuths indicated by the arrow heads.


tude. To correlate these values in various sections of the country, the values in seconds of arc were reduced to linear measure with 10 feet as the unit. This unit gave the best balance for these quantities in the formation of the normal equations. In this manner the set of equations for latitude and the set for longitude shown in the table on page 16 were formed. The constant terms in these equations must be multiplied by 10 if it is desired to get the closures in feet.

After these two sets of observation equations have been formed, it becomes necessary to decide upon a proper weighting of the various sections. If they were all of the same length, the weight of each sec-



tion could be taken as unity, but such a condition would probably never be found in actual practice. In this adjustment it was decided to weight the sections inversely proportional to their length. A sec-

tion of about average length was assumed as having weight unity. A section approximately one-half as long would then have the weight



2, while a section twice as long as the unit section would have the weight 0.5, and so on. No attempt was made to get these relative lengths with any great exactness; they were actually scaled roughly

from a progress map, since a slight variation of the weight would have but little effect on the final value of the resulting v for the given section.

In this way there was formed a set of 42 observation equations for latitude and the same number for longitude, this being the number of sections in the whole net. The symbols x_1, x_2 , etc., were used as corrections to the assumed latitudes of the junction points, and y_1, y_2 , etc., were used as corrections to the assumed longitudes of the same points. The 42 expressions for the v's gave 26 normal equations in each case, this being the number of the junction points, omitting, of course, Meade's Ranch which was adopted as the fundamental position and which was not to change from the assumed value.

After the solution of the normal equations was made and the values of the x's and y's were determined, the expressions in units of 10 feet were reduced back to fractions of a second of latitude and longitude. When these values were added to the assumed latitudes and longitudes the position of a junction point in each junction figure was determined. A correction was then made to the Laplace azimuth or to the adopted azimuth as the case might be, due to the change in the geodetic longitude in the junction figure. After this was done all of the positions of the stations in the junction figures were computed. This gave a line at each end of the various sections that was fixed in all respects—in length, in azimuth, and in position upon the ellipsoid of reference. The sections could then be adjusted with inclusion of the equations for closure in latitude and longitude and thus the whole net could be placed on one continuous consistent datum.

FORMATION OF OBSERVATION EQUATIONS

The table of the observation equations on page 16 makes it practically self-evident as to how the equations are formed. However, to show the method in cases where there is found a failure in closure, we will give a sample of the formation of one such in the list. Between Donna and Peters (see tables on pp. 14 and 15) we have the following differences:

]	n latitude		I	n longitude
Assumed difference	3 3	, 00 00	52. 382-x ₄ +x ₅ 52. 154+v ₅	° 2 2	, 33 33	17. $613 - y_4 + y_5$ 17. $712 + v_5$

Since, after adjustment, these values should agree, we find for latitude that $v_5 = +0.228 - x_4 + x_5$, and for longitude that $v_5 = -0.099 - 0.000 - 0.000 - 0.000 - 0.000 - 0.000 - 0.000 - 0.000 - 0.000 - 0.00$

This discrepancy of closure was developed at station Peters and at this latitude 0.228 in latitude is equal to 23.0 feet and 0.099 in longitude is equal to 8.8 feet. For convenience in the solution, as stated on page 8, the unit was assumed as 10 feet; accordingly, the observation equations become, in latitude

$$v_5 = +2.30 - x_4 + x_5$$

and in longitude

$$v_5 = -0.88 - y_4 + y_5$$

In the same way all of the other observation equations were formed.

From the procedure indicated above, it is seen that the extent in latitude and longitude of each section is considered as if it were a quantity directly measured; the v's for each section are the corrections in latitude and in longitude that are to be applied to these quantities. The solution of the observation equations by least squares gives in each case the most probable distribution of the closures at the junction points. The later adjustment of the sections between the junction figures, thus fixed in all respects upon the spheroid (that is, fixed in length, position, and azimuth) brings about a just distribution of the position closures among the various are sections in the whole net of triangulation.




FIGURE 6.-Section closures resulting from apportionment of loop closures

The first number above the line is the closure in latitude in feet, the second number the closure in longitude in feet, and the third number the total closure in feet. On the other side of the line representing the section is the approximate proportional part of the length of the section represented by the total closure.

SOLUTION OF NORMAL EQUATIONS

The forward solution of the normal equations can be considerably shortened by carrying the solution of the latitude equations and that of the longitude equations at the same time. The coefficients of the x's are the same as the coefficients of the corresponding y's, and hence the only differences between the two solutions are in the η and Σ columns. These can both be appended to the one solution, as is done in the solution given on pages 21 to 25 and much computation is avoided in this way. The control of the solution of each set is given in its respective Σ column, and hence the single solution is just

as effective as a double solution would be. As a further control of the solution, the values of $(p\eta^2)$, $(p\eta\Sigma)$ and $(p\Sigma^2)$ are appended to the normals and the resulting work is appended to the forward solution. (See p. 25.) It should be noted that the values in the columns η and Σ on page 25 for the latitude equations are derived in the regular way, and that the corresponding values in the longitude set are similarly derived just as if the latitude values were not present. In other words, the cross multiplication for the latitude set is only applied to the η and Σ column of that set and then the corresponding multiplier is used for the longitude set. After adding up the three columns occurring beyond the solution of equation No. 26, the three results




Figure 7.—Changes in positions of junction points, previously adjusted; caused by the readjustment

The sign indicates in which direction the position was changed. The upper number is the change in latitude and the lower number the change in longitude.

should be equal, each being the sum of the respective pv^2 's. It will be seen that with the exception of the unavoidable uncertainty in the last places of decimals, the same result is obtained in both sets of normals. It will also be seen from the sum of the pv^2 's given in the table on page 28 that these solution values are checked except for the uncertainty due to the dropping of decimals.

COMPUTATIONS

On the following pages are assembled in tabular form the computations involved in adjusting the latitude and longitude closures of the junction points.

U. S. COAST AND GEODETIC SURVEY

Assumed positions of junction stations

Station	As	sum	ed latitude	Ass	ıme	d longitude
	•	,	"	۰		"
Carson	35	16	24. $889+x_1$	97	57	32. $437+y_1$
Kyle	32	49	18. $024 + x_2$	98	19	11. $654+y_2$
Mission.	29	42	52. $781+x_3$	98	09	$52.133+y_3$
Donna	26	09	40. 491 $+x_4$	98	02	$44.447 + y_4$
Peters	29	10	32.873 $+x_5$	100	36	$02.060+y_5$
Ingle	31	35	48.698+x6	103	35	24. 147+y ₆
Phillips	34	59	03. $367 + x_7$	104	07	$59.481 + y_7$
Aroya	38	48	09. $701+x_8$	103	10	$55.504+y_8$
Whitetank	33	34	01. $744+x_9$	112	33	$27.214+y_9$
Mount Nebo	39	48	38. $324+x_{10}$	111	45	$56.284+y_{10}$
Black	35	55	$51.691+x_{11}$	115	02	$35.657 + y_{11}$
Mount Diablo	37	52	54. 246 $+x_{12}$	121	54	
Tepusquet	34	54	36. 391 $+x_{13}$	120	11	$08.121+y_{13}$
Farmer	43	42	21. 280+ x_{14}	97	40,	
States	49	00	01. 298 $+x_{15}$	97	07	$39.003 + y_{15}$
Sundance	44	28	44. $100+x_{16}$	104	. 27	02. 892+y ₁₆
Noige	48	53	37. $986+x_{17}$. 103	47	$20.841+y_{17}$
Oxford	42	16	10.863 $+x_{18}$	112	05	49. 476 $+y_{18}$
Mount Ellis	45	34	38. 959+ x_{19}	110	57	$18.429+y_{19}$
Goldstone	48	52	55. $718+x_{20}$	110	29	18. $696 + y_{20}$
Dry	44	10	$08.208+x_{21}$	117	39	$39.828+y_{21}$
Alder	45	50	$59.400+x_{22}$	119	56	$21.043+y_{22}$
Bally	40	36	10. $730+x_{23}$	122	38	$59.025 + y_{23}$
Red		56	$06.077 + x_{24}$	121	49	$11.048 + y_{24}$
Oroville	48	53	44. $253+x_{25}$	119	20	12. $795+y_{25}$
Birch Point	48	56	30. $780+x_{26}$	122	49	12. $025+y_{26}$

Fixed position 1

Station		Lati	tude	I	ong	itude
Meade's Ranch	39	, 13	26. 686	98	, 32	30. 506

¹ This is the fixed position on which the North American Datum depends.

Computed differences of positions of junction stations

1 Carson-Meade's Ranch -3 57 01,797 -0 34 58,069 28 Kyle-Carson -2 27 06,865 +0 21 39,217 30 Mission-Kyle -3 06 25,243 -0 09 19,521 4 Donna-Mission -3 33 12,290 -0 07 07,686 5 Peters-Donna +3 00 52 154 +2 33 17,712 6 Peters-Mission -0 32 20,068 +2 26 10,142 7 Ingle-Peters +2 25 15,825 +2 59 22,087 8 Ingle-Kyle -1 13 29,224 +5 16 12,826 9 Phillips-Carson -0 17 21,648 46 10 27,188 10 Aroya-Meade's Ranch -0 25 16,985 +4 38 24,998 11 Phillips-Ingle +3 23 14,669 +0 32 35,334 12 Aroya-Phillips +3 49 06,334 -0 57 03,977 13 Whitetank-Ingle +1 31 30,466 +8 38 30,667 14 Whitetank-Phillips -1 25 01,810 +8 25 27,838 15 Mount Nebo -7 35 246,633 +3 16 39,373 17 Black-Mount Nebo -3 52 46,633 +3 16 39,373 17 Black-Whitetank +2 21 50,021 +2 29 08,021 18 Mount Diablo-Mount Nebo -1 55 43,917 +10 08 50,926 19 Tepusquet-Mount Diablo -2 58 17,855 +7 37 40,852 21 Farmer-Meade's Ranch +4 28 54,594 +10 08 50,926 22 States-Farmer +0 46 23,031 +6 46 29,153 24 Sundance-Farmer +0 46 23,031 +6 46 29,153 24 Sundance-Farmer +0 46 23,031 +6 46 29,153 24 Sundance-Aroya +5 40 34,399 +1 16 67,388 27 Oxford-Mount Nebo +2 27 32,539 +0 19 53,192 28 Mount Ellis-Oxford +3 18 28,096 -1 08 31,047 47,977 36 Bally-Mount Diablo +2 43 16,484 +0 44 12,113 46 46 47,977 47,977 48 Bally-Mount Diablo +2 43 16,484 +0 44 12,113 48 49 40 47,977 41 40 51,192 +2 46 41 51,192 +2 45 45 45 45 45 45 45 4	No. of differ- ence	Junction stations	Diffe	rene tu	e of lati- le		enc gitu	e of lon-
1				,		•		. ,,
2 Kyle-Carson	1 ,	Carson-Meade's Ranch	_					
3 Mission-Kyle	2	Kyle-Carson	2	27	06.865	+0	21	39. 217
5 Peters-Donna +3 00 52.154 +2 33 17.712 6 Peters-Mission -0 32 20.068 +2 28 10.122 8 Ingle-Peters +2 25 15.825 +2 59 20.2087 8 Ingle-Kyle -1 13 29.224 +5 16 12.826 9 Phillips-Carson -0 17 21.648 +6 10 27.828 10 Aroya-Meade's Ranch -0 25 16.985 +4 38 24.998 11 Phillips-Ingle +3 49 06.334 -0 57 03.977 14 Whitetank-Ingle +1 58 13.046 +8 88 03.097 14 Whitetank-Phillips -1 25 01.810 +8 25 27.838 15 Mount Nebo -3 52 46.633 +3 16 39.273 16 Black-Mount Nebo </td <td></td> <td>Mission-Kyle</td> <td>· -3</td> <td></td> <td></td> <td></td> <td></td> <td></td>		Mission-Kyle	· -3					
Topic February Topic Topi		Peters-Donna	+3					
Red-Ayle	6	Peters-Mission	-0					
9 Phillips-Carson		Ingle-Peters	+2	25.				
10	ŝ	Phillips-Carson	-0					
12		Aroya-Meade's Ranch	-ŏ					
13		Phillips-Ingle	+3					
14		Whitetank-Ingle	1 1					
15 Mount Nebo-Aroya	14	Whitetank-Phillips	-1	25	01.810	+8	25	27.838
Black-Whitetank	15	Mount Nebo-Aroya	+1	00	28. 623	+8	35	00. 780
18		Black-Mount Nebo	-3					
Tepusquet-Mount Diablo								
Tepusquet-Whitetank								
22	20	Tepusquet-Whitetank	+ī	20	34.895	+7	37	
23		Farmer-Meade's Ranch				-0		56. 400
24		States-Farmer	+5					
25		Sundance-Arova	+5					
27	25	Norge-Sundance	+4					
28	26	Norge-States	-0					
29 Mount Ellis-Sundance		Oxford-Mount Nebo	+2					
30 Goldstone-Mount Ellis	20	Mount Ellis-Sundance	±3					
32 Dry-Oxford	30	Goldstone-Mount Ellis	+3					
33 Alder-Dry	31	Goldstone-Norge	-0					
34 Red-Alder								
35 Red-Bally		Red-Alder	ΙĮά					
37 Bally-Dry		Red-Bally	+5					
38		Bally-Mount Diablo	+2					
39 Oroville-Alder		Bany-Dry	-3					
40 Tepusquet-Black ————————————————————————————————————		Oroville-Alder	+3					
41 Birch Point-Oroville		Tepusquet-Black	-ī					
42 Birch Point-Red		Birch Point-Oroville						
	42	Birch Point-Red	+3	00	24.703	+1	00	00. 977

Observation equations

Latitude equation	Weight,	Longitude equation
$\begin{array}{c} v_1 = x_1 \\ v_2 = x_2 - x_1 \\ v_3 = x_3 - x_2 \\ v_4 = x_4 - x_3 \\ v_5 = +2, 30 - x_4 + x_5 \end{array}$	1. 00 1. 65 1. 45 1. 00 . 90	$v_1 = y_1 v_2 = y_2 - y_1 v_3 = y_3 - y_2 v_4 = y_4 - y_3 v_5 = -0.88 - y_4 + y_5$
$\begin{array}{c} v_6 = +1.62 - x_3 + x_5 \dots \\ v_7 = x_6 - x_5 \dots \\ v_8 = -1.03 - x_2 + x_6 \dots \\ v_9 = +1.27 - x_1 + x_7 \dots \\ v_{10} = x_8 \dots \end{array}$	1. 65 1. 00 . 75 . 70 1. 10	$v_6 = -1.91 - y_3 + y_5$ $v_7 = y_6 - y_5$ $v_8 = -2.88 - y_2 + y_6$ $v_9 = -1.20 - y_1 + y_7$ $v_{10} = y_8$
$\begin{array}{c} v_{11} = x_7 - x_5 \\ v_{12} = x_8 - x_7 \\ v_{13} = x_9 - x_6 \\ v_{14} = +1.89 - x_7 + x_9 \\ v_{15} = x_{10} - x_8 \end{array}$	1. 10 1. 00 . 50 . 50 . 70	$\begin{array}{l} v_{11} = y_7 - y_6 \\ v_{12} = y_8 - y_7 \\ v_{13} = y_9 - y_6 \\ v_{14} = -0.89 - y_7 + y_9 \\ v_{15} = y_{10} - y_8 \end{array}$
$\begin{array}{c} v_{16} = x_{11} - x_{10} \\ v_{17} = -0.75 - x_9 + x_{11} \\ v_{18} = -1.63 - x_{10} + x_{12} \\ v_{19} = x_{13} - x_{12} \\ v_{20} = -2.51 - x_9 + x_{13} \\ \end{array}$	2. 00 2. 00 . 70 2. 00 . 90	$\begin{array}{l} v_{16} = y_{11} - y_{10} \\ v_{17} = +3.57 - y_{9} + y_{11} \\ v_{18} = -2.39 - y_{10} + y_{12} \\ v_{19} = y_{13} - y_{12} \\ v_{20} = +0.47 - y_{9} + y_{13} \end{array}$
$\begin{array}{c} v_{21} = x_{14} \\ v_{22} = -2.55 - x_{14} + x_{15} \\ v_{23} = -2.14 - x_{14} + x_{16} \\ v_{24} = x_{16} - x_{8} \\ v_{25} = x_{17} - x_{16} \end{array}$. 80 . 65 . 80 . 80	$\begin{array}{l} v_{21} = y_{14} \\ v_{22} = -2.91 - y_{14} + y_{15} \\ v_{23} = -2.70 - y_{14} + y_{16} \\ v_{24} = y_{16} - y_{8} \\ v_{25} = y_{17} - y_{16} \end{array}$
$\begin{array}{c} v_{26} \! = \! x_{17} \! - \! x_{15} \\ v_{27} \! = \! x_{18} \! - \! x_{10} \\ v_{28} \! = \! x_{19} \! - \! x_{18} \\ v_{29} \! = \! + \! 1.58 \! - \! x_{16} \! + \! x_{19} \\ v_{30} \! = \! x_{20} \! - \! x_{19} \end{array}$. 85 2. 00 1. 65 . 70 1. 25	$\begin{array}{c} v_{26} = y_{17} - y_{15} \\ v_{27} = y_{18} - y_{10} \\ v_{28} = y_{19} - y_{18} \\ v_{29} = +0.97 - y_{16} + y_{19} \\ v_{30} = y_{20} - y_{19} \end{array}$
$\begin{array}{c} v_{31} = +1.56 - x_{17} + x_{20} \dots \\ v_{32} = x_{21} - x_{18} \dots \\ v_{33} = x_{22} - x_{21} \dots \\ v_{34} = x_{24} - x_{22} \dots \\ v_{35} = x_{24} - x_{23} \dots \end{array}$. 80 1. 25 2. 00 2. 00 . 90	$v_{31} = +3.05 - y_{17} + y_{20}$ $v_{32} = y_{21} - y_{18}$ $v_{33} = y_{22} - y_{21}$ $v_{34} = y_{24} - y_{22}$ $v_{35} = y_{24} - y_{23}$
$ \begin{vmatrix} v_{36} = x_{23} - x_{12}, \\ v_{37} = -2. \ 91 - x_{21} + x_{23}, \\ v_{38} = x_{25} - x_{20}, \\ v_{39} = +0. \ 66 - x_{22} + x_{25}, \\ v_{40} = -0. \ 77 - x_{11} + x_{13}, \\ \end{vmatrix} $	2.00 .80 .70 1.20 .80	$\begin{array}{l} v_{36} = y_{23} - y_{12} \\ v_{37} = -1.03 - y_{21} + y_{23} \\ v_{38} = y_{25} - y_{20} \\ v_{39} = -0.25 - y_{22} + y_{25} \\ v_{40} = -2.95 - y_{11} + y_{13} \end{array}$
$v_{41} = -1.35 - x_{25} + x_{26} - \dots $ $v_{42} = x_{26} - x_{24} - \dots $	1. 45 1. 25	$v_{41} = +0.94 - y_{25} + y_{26} v_{42} = y_{26} - y_{24}$

Table for formation of latitude normal equations

р		7	1	2	3	4	- 5	6	7	8	. 8	10	11	12	13	14	15	16	17	18	19	20	21	22	23 .	24	25	26	рη	ρΣ
1.00	5 -	0	+1																										0	+1.00
1. 68		0	-1	+1													1												O	0
1.45	5	0		-1	+1			ļ										J											O O	0
1.00		0 20			-1	+1																							0 +2.0700	0 +2.07
	١.	⊦2.30				-1	+1										i			ļ										
1.65		-1.62			-1		+1						 -				[ļ			[-		+2.6730	+2.6
1.00		9					-1	+1 +1													ļ								0 7725	0 7
. 75	3 7	-1. 03 -1. 27		-1				+1																					+. 8890	+.8
1.10	₹∥₹	0 Z	-1						+1	+1													-						7.∞∞	+1.1
		Ť								41											 									1
1. 10	? }	0				·;		-1	+1								[} <i>:</i>						¦		¦			Ŏ	į
1.00 .50		8						:-	-1	+1																-			0	l ö
50		-1.89						-1			\+1 +1																		+. 9450	+.9
1 :70	ין ג	0							-1		1	+1																	0	0.
1		·]								_1		١.								}									Ô	1 .
2.00		0							[:-	-1	+1					-				[-1. 5000	-1.5
2,00		75 -1. 63									-1	;-	+1	,-;-															-1.1410	-i. i
2.00	(I-	0										-1		+ 1	+1			}									-		0 1410	0
2.90		-2.51									-1			ļ —.	Ŧi														-2. 2590	-2.2
		1							1						1 7.														0	+.8
. 80	?	0 -2.55														+ <u>1</u>	,-;-												-1. 6575	-1.6
80	: II	-2. 14														-1	+1	+1											-1.7120	-i.
.80	i l'	0 1								-1						-1		Ŧi											0.1120	ا ق
80		ŏ																<u>-</u> î	+1										Ŏ	ŏ
. 85	- 1	0				1	1	1	1					1			1	1 -		1	1				1				^	۱۸
2.00		ŏ]				<u>-</u> -					-1		+1	+1									Ä	X
1.65		ŏ										- 1								Ti	+1								ŏ	lŏ
7.70	í l∓	-i. 58																-1		_*	∓î								+ĭ. 1060	+1.1
1. 25	5 I '	ō															1				l –ī	+1	1						0	0
. 80	- 1	-1. 58				1	1	1								1	1	1	-1	1	1 -	+1	1	1			1	1 !	+1.2480	+1.2
1. 25	ין י	0.00					[-			-1			┯∗	+1	ļ					0	1 0
2.00		ŏ							1											l*	1		-î	+1					Ŏ	Ŏ
2.00) l	ŏ											l							<u> </u>			<u> </u>	<u>-</u> î		+1			Ō	ĬŎ
. 90		ŏ Ì										l			l										-1	+1		l	0	0
2.00		0		,					1			}		_1	1		1	1	1				1	1	+1	lí	1		0	0
.80	۱L_	-2. 91												_,		··									l ∓i		1		-2.3280	-2.3
.70	۱ آ	0				1	l		1					<u> </u>	1		1					-1	l		1		+1	1	Ō	0
1. 20	١.	+. 66							1						I							- -		-1		1	+1		+. 7920	+.
. 80		77											-1		+1			ļ			.]			ļ	ļ				6160	(
1.45		-1. 35		ļ.,	1	1	t i	1	1		1	l	1	1	l .	1	1	1	l	ļ	1	1	1	1	Į.		1 -1	+1	-1.9575	-1.8
1. 25		0												1				1	ļ		1						1	⊢i	0.0010	, î

Table for formation of longitude normal equations—Continued

0	p	η	1	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	p_{η}	pΣ .
	. 00 . 65 . 45	0	•	+1 -1	+1 -1	+1																								0 0	+1.00 0 0
5 6 1	. 00 . 90 .:65	0 -1.				-1 -1	+1 -1	+1 +1																						0 7920 -3. 1515	0 7920 -3. 1515
	. 75 . 70 . 10	0 -2. -1. 0		-1	-i			-i	+1 +1	+1	+1																			0 -2. 1600 8400	0 -2.1600 8400 +1.10
11 1 12 1 13	. 10 . 00 . 50	0							-1 -1	+1 -1	+1	+1																		0	0 0
16 2	. 50 . 70 2. 00	0								—1	- <u>1</u>	+1	+1 -1	+1																4450 0 0	4450 0 0
18 19 2	2.00 .70 2.00 .90	+3. -2. 0 +.	39									-1 1	-1	+1	+1 -1	+1 +1			4						 					+7. 1400 -1. 6730 0 +. 4230	+7. 1400 -1. 6730 0 +. 4230
22 23	. 80 . 65 . 80	0 -2. -2.	91														+1 -1 -1	+1	+1											0 -1.8915 -2.1600	+.80 -1.8915 -2.1600
	. 80 . 80 . 85	0									—1 							1	+1 -1	+1 +1										0	0 0 0
28 1 29 1	2.00 .65 .70	0+0											-1 						 1		+1 1	+1 +1 -1	+1							0 0 + 6790	0 0 +. 6790
31 32 1 33 2	. 80 . 25 2. 00	+3. 0																		-1	-i		+1	+1 -1	+i					+2.4400 0	+2. 4400 0
34 2 35 36 2	2.00 .90 2.00	0													-1										-1	-1 +1	+1 +1			0	0
39 1	. 80 . 70 . 20 . 80	-1. 0 -2.	25													+1							-i	-1	<u></u> 1	+1 		+1 +1		8240 0 3000 -2. 3600	8240 0 3000 -2. 3600
- 1	. 45 . 25	+. 0																									-1	-1	+1 +1	+1. 3630 0	+1. 3630 0

Normal equations for latitude

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16 17	18	η	Σ
1 2 3 4 5	+3.35	-1. 65 +3. 85	-1.45 +4.10	-1.00 +1.90	-1.65 90	-0.75			- -									+. 7725 -2. 6730	+0. 1110 +. 7725 -2. 6730 -2. 0700
5 6 7 8 9				+1.90	+3.55	-1.00 +3.35			-0. 50 -, 50							0.80		+4. 7430 7725	+4. 7430 7725
10	11	12	13	14		15	16	17	18	+5. 40 19	20	-0. 70 21	22	23	24	25	26	7 +1. 1410	+1. 1410 E
11 12 13 14 15	+4. 80	+4.70	-0. 0 -2. +3.	00											0			-0. 8840 -1. 1410 -2. 8750	-0. 8840 -1. 1410 -2. 8750
1			+0.	+2		1, 50		-0.85				-						+3. 3695 -1. 6575	+4. 1695 -1. 6575 -2. 8180
16 17 18 19 20								+2.45	+4.90	-0.70 -1.65 +3.60	-0.80 -1.25 +2.75	-1.2	5			-0.70		-1. 2480 0 +1. 1060 +1. 2480	-2.8180 -1.2480 0 +1.1060 +1.2480
21 22 23 24 25												+4.0	-2. +5.	008 20 +3.7	-2.0	Ď	-1. 25 -1. 45	+2.3280 7920 -2.3280 0 +2.7495	+2.3280 7920 -2.3280 0 +2.7495
26					-												+2.70	-1.9575 +43.4557	-1. 9575

Normal equations for longitude—Continued

	1	2	3	4	5	6	-7	8	9	10	11	12	13	14	15	16	17	18	ŋ	Σ
1 2 3 4 5 6 7 8 9	-	-1.65 +3.85	-1.45 +4.10	-1.00 +1.90	-1. 65 90 +3. 55	-0. 75 -1. 00 +3. 35	-		-0. 50 50		-2.00					-0.80			+3. 1515 +. 7920 -3. 9435 -2. 1600 3950	-3. 9435 -2. 1600
	11	12	13	14	1	15	16	17	18	19	20	21	22	23	3 :	24	25	26	η	Σ
11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26		+4.70	-0.5 -2.6 +3.	70	. 25 -+	1. 50						-1. 2	6	-2	. 80	2.00 90 4.15	-0. 70 -1. 20	—1. 25 —1. 45	+9. 5000 -1. 6730 -1. 9370 +4. 0515 -1. 8915 -2. 4390 -2. 4400 +. 6790 +2. 4400 +. 8240 +. 3240 -1. 6630 -1. 6630 +72. 6321	+9.5000 -1.6730 -1.9370 +4.8515 -1.8915 -2.8390 -2.4400 0 +6790 +2.4400 +8240 0 -5240 0 -1.6630 +1.3630 +72.6321 +75.5321

Solution of normal equations

1	2	3	4	5	6	7	8	9	η (Lat.)	Σ (Lat.)	η (Long.)	Σ (Long.)
+3.35 21=	-1.65 +.49254					-0.70 +.20896			-0.8890 +.26537	+0.1110 03313	+0. 8400 25075	+1.8400 54925
1	+3.85 8127	-1.45			-0.75	3448			+. 7725 4379	+. 7725 +. 0547	+2 1600 +. 4137	+2.1600 +.9063
	$+3.0373$ $x_1 =$	-1.45 +.47740			75 +. 24693	3448 +. 11352			+. 3346 11016	+. 8272 1 27231	+2.5737 84736	+3.0663 2 -1.00952
	2	+4. 10 6922	-1.00	-1.65	3580	 1646			-2.6730 +.1597	-2.6730 +.3949	+3. 1515 +1. 2287	+3. 1515 +1. 4638
		+3. 4078 x ₁ =	-1.00 +.29344	-1.65 +.48418	3580 +. 10505	1646 +. 04830			-2.5133 +.73751	-2. 2781 +. 66850	+4.3802 -1.28535	+4. 6153 4 -1. 35436
		3	+1. 90 2934	90 4842	1050	0483			-2.0700 7375	-2.0700 6685	+. 7920 +1. 2853	+. 7920 +1. 3543
			+1.6066 x ₄ =	-1.3842 .+.86157	1050 +. 06536	0483 +. 03006		·	-2.8075 +1.74748	-2.7385 4 +1.70447	+2.0773 -1.29298	+2.1463 4 -1.33599
			3	+3. 55 7989 -1. 1926	-1.00 1733 0905	0797 0416			+4.7430 -1.2169 -2.4189	+4. 7430 -1. 1030 -2. 3593	-3. 9435 +2. 1208 +1. 7897	-3. 9435 +2. 2347 +1. 8493
}				+1. 5585 z ₆ =	-1. 2638 +. 81091	1213 +. 07783			+1. 1072 71043	+1. 2807 6 82169	0330 +. 02017	+ 1405 4 - 09009
				2 3 4 5	+3.35 1852 0376 0069 -1.0248	-1. 10 0851 0173 0032 0984		-0.50	7725 +. 0826 2640 1835 +. 8978	ー. 7725 十. 2042 ー. 2393 ー. 1790 十1. 0385	-2. 1600 +. 6355 +. 4601 +. 1358 0268	-2. 1600 +. 7571 +. 4848 +. 1403 +. 1139
ļ					+2.0955 z ₀ =	-1.3040 +.62229		50 +. 23861	2396 +. 11434	+. 0519 02477	9554 +. 45593	6639 +. 31682
					1 2 3	+3.30 1463 0391	-1.00		0560 1858 +. 0380	0560 +. 0233 +. 0939	3950 +. 1755 +. 2922	3950 +. 3845 +. 3481 +. 2229
					3 4 5 8	0080 0015 0094 8115		 3111	1214 0844 +. 0862 1491	1100 0623 +. 0997 +. 0323	+. 2116 +. 0624 0026 5945	+. 2229 +. 0645 +. 0109 4131
						+2. 2842 z ₇ =	-1. 00 +. 43779	8111 +. 35509	4725 +. 20686	+. 0006 00026	2504 +. 10962	+. 2228 7 09750

	8	9	10	12	14	15	16	17	18	19	23	11	13	η (Lat.)	Σ (Lat.)	η (Long.)	Σ (Long.)
7	+3.60 4378	3551	-0.70				-0.80							0 2069	+1.10 +.0003	0 1096	+1.10 +.0975
	+3. 1622 x ₈ =	3551 +. 11230					80 +. 25299				 			2069 +. 06543	+1.1003 2 34792	1096 +. 03466	+1. 1975 37869
	6 7	+3. 90 1193 2880 0399					••••					-2.00	-0.90	+4. 7040 0572 1678 0232	+4.7040 +.0124 +.0002 +.1236	-8.0080 2280 0889 0123	-8. 0080 1584 +. 0791 +. 1345
	Ů	+3. 4528 x ₀ =	0786 +. 02276				0898 +. 02601	 				-2. 00 +. 57924	90 +. 26066	+4. 4558 -1. 29049	+4.8402 -1.40182	-8.3372 +2.41462	-7. 9528 +2. 30329
		8 9	+5. 40 1550 0018				1771 0020					-2.00 0455	0 2 05	+1. 1410 0458 +. 1014	+1. 1410 +. 2435 +. 1102	+1.6730 0243 1898	+1.6730 +.2651 1810
			+5. 2432 x ₁₀ =	70 +. 13351 +4. 70			1791 +. 03416		-2.00 +.38145		-2.00	-2. 0455 +. 39012	0205 +. 00391 -2. 00	+1. 1966 22822 -1. 1410	+1. 4947 28507 -1. 1410	+1. 4589 27825 -1. 6730	+1.7571 0 33510 -1.6730
			10	0935 +4. 6065			0239		2670 2670		-2.00	2731 2731	0027 -2. 0027	+. 1593	+. 1990 9420 19	+. 1943	+. 2339 -1. 4391 89
	i			$x_{12} =$	+2. 25 x ₁₄ =	-0. 65 +. 28889	80				+. 43417			+. 21311 +3. 3695 -1. 49756	+. 20447 +4. 1695 -1. 85311	+. 32100 +4. 0515 -1. 80067	+. 31236 +4. 8515 -2. 15622
	<u> </u> !		·		14	+1.50	- 2311	-0.85						-1. 6575 +. 9734	-1. 6575 +1. 2045	-1.8915 +1.1704	-1.8915 +1.4015
					,	+1.3122 x ₁₅ =								6841 +. 52134	4530 +. 34522	7211 +. 54954	4900 +. 37342
						8 9 10	0023 0061		 0683	-0. 70		0520 0699	0234 0007	-2.8180 0523 +.1159 +.0409	-2.8180 +.2783 +.1259 +.0511	-2. 8390 0277 2169 +. 0498	-2.8390 +.3030 2069 +.0600
			-			12 14 15	—. 2844	1497				0014	0104	0051 +1. 1981 1205	0049 +1. 4825 0798	0077 +1. 4406 1270	0075 +1. 7250 0863
							$+2.5640$ $x_{16} =$	一. 9497 十. 37040	0697 +. 02718	70 +. 27301	0104 +. 00406			-1.6410 +.64002	9649 6 +. 37621	-1.7279 +.67391	-1.0517 5 +.41010

	17	18	19	20	21	22	23	11	13	24	25	η (Lat.)	Σ (Lat.)	η (Long.)	Σ (Long.)
15 16	+2.45 5506 3518	-0. 0258	-0. 2593	-0.80			-0.0039	-0.0457	-0.0128			-1. 2480. 4431 6078	-1. 2480 2934 3573	-2. 4400 4671 6400	-2. 4400 3174 3895
	$+1.5476$ $x_{17}=$	0258 +. 01667	2593 +. 16755	80 +. 51693			0039 +. 00252	0457 +. 02953	0128 +. 00827			-2. 2989 +1. 48546	-1.8987 8 +1.22693	-3. 5471 +2. 29200	-3. 1469 70 +2. 03347
	10 12 16 17	+4. 90 7629 0155 0019 0004	-1. 65 0190 0043	—. 0133	-1. 25		1159 0003 0001	7803 0158 0034 0008	0078 1161 0009 0002			0 +. 4564 0569 0446 0383	0 +. 5702 0546 0262 0317	0 +. 5565 0857 0470 0591	0 +. 6702 0834 0286 0525
		+4. 1193 x ₁₈ =	-1. 6733 +. 40621	0133 +. 00323	-1. 25 +. 30345		1163 +. 02823	8003 +. 19428	1250 +. 03034			+. 3166 07686	+. 4577 11111	+. 3647 08853	+. 5057 8 12279
		16 17 18	+3. 60 1911 0434 6797	-1. 25 1340 0054	5078		0028 0007 0472	0337 0077 3251	0094 0021 0508		·	+1. 1060 4480 3852 +. 1286	+1.1060 - 2633 - 3181 +.1859	+. 6790 4717 5943 +. 1481	+ 6790 - 2871 - 5273 + 2055
			+2.6858 x ₁₀ =	-1.3894 +.51731	5078 +. 18907		一. 0507 十. 01888	3665 +. 13646	0623 +. 02320			+. 4014 14945	+. 7105 26454	2389 +. 08895	+. 0701 2 02614
			17 18 19	+2.75 4135 0 7188	0040 2627		0020 0004 0262	0236 0026 1896	0066 0004 0322		-0.70	+1. 2480 -1. 1884 ·+. 0010 +. 2076	+1. 2480 9815 +. 0015 +. 3675	+2.4400 -1.8336 +.0012 1236	+2. 4400 -1. 6268 +. 0016 +. 0363
				+1.6177 120==	2667 +. 16486		一. 0286 十. 01768	2158 +. 13340	0392 +. 02423		70 +. 43271	+. 2682 16579	+. 6355 6 39290	+. 4840 29919	+. 8511 4 52630
				18 19 20	+4. 05 3793 0960 0440	-2.00	80 0353 0096 0047	2429 0693 0356	0379 0118 0065		—. 1154	+2.3280 +.0961 +.0759 +.0442	+2.3280 +.1389 +.1343 +.1048	+. 8240 +. 1107 0452 +. 0798	+. 8240 +. 1535 +. 0133 +. 1404
					$+3.5307$ $x_{21} =$	-2.00 +.56646	8496 +. 24063	3478 +. 09851	0562 +. 01592		1154 +. 03268	+2.5442 72059	+2.706Ø 59 76639	—. 27453	+1. 1312 0 - 32033
					21	+5. 20 -1. 1329	4813	1970	0318	-2.00	-1. 20 0654	7920 +1. 4412	7920 +1. 2328	+. 3000 +. 5491	+. 3000 +. 6407
						+4.0671 x ₂₂ =	4813 +. 11834	1970 +. 04844	0318 +. 00782	-2.00 +.49175	1. 2654 +. 31113	+. 6492 15962	+. 7408 18214	+. 8491 20877	+. 9407 23130

1					 				,
	23	11	13	24	25	η (Lat.)	Σ (Lat.)	η (Long.)	Σ (Long.)
12 16 17 18 19 20 21 22	+3.708683 0 000330010000520440570 +2.5655 x22= 9 10 112 16 17 18 19 20 21 222	-0. 1186 0005 0001 0226 0069 0038 0837 0233 2595 +. 10115 +4. 80 1. 1585 7980 0162 0059 0038 0590 0288 0343 0099	-0. 8695 0001 0035 0012 0037 0135 0038 8923 +. 34781 89 5213 0080 1187 0017 0004 0243 0085 0052 0055 0015	0969	0934 0114 0613	-2. 3280 4262 0067 0058 +. 0089 +. 0076 +. 0047 +. 6122 +. 0768 -2. 0565 +. 80160 8840 +2. 5810 +. 4668 0582 0789 +. 0615 +. 0548 +. 0358 +.	-2. 3280 4089 0039 0048 +. 0129 +. 0134 +. 0112 +. 6511 +. 0877 -1. 9693 4 +. 76765 8840 +2. 8036 +. 5831 0464 +. 0859 0464 +. 0866 +. 0848 +. 0848	-0. 8240 6420 0070 0089 +. 0103 0045 +. 2332 +. 1005 -1. 1338 +. 44194 +9. 5000 -4. 8292 +. 5691 0871 0831 +. 0708 0326 +. 0646 0955 +. 0411	-0. 8240 -0. 6247 0043 0079 +. 0143 +. 0151 +. 2722 +. 1113 -1. 0467 +. 4. 4079 +9. 5000 -4. 6066 +. 6854 0823 0929 +. 0983 +. 1136 +. 1136 +. 1136 +. 1136 +. 1144 +. 0456
	23	0262 $+2.5158$ $x_{11}=$	0903 -1. 5854 +. 63018	- 1150 - 2119 + 08423	0192 1853 + 07365	2080 +2. 1849 86847	1992 +2. 7181 -1. 08041	1147 +5. 0891 -2. 02286	1059 +5. 6226 3 -2. 23480
		9 10 12 16 17 18 19 20 21 22 23 11	+3. 70 2346 0001 8707 0005 0001 0038 0014 0009 0009 0002 3104 9991 +1. 2773			-2.8750 +1.1614 +.0047 4268 0221 0190 +.0093 +.0065 +.0405 +.0051 7153 +1.3769 -1.4442	-2 8750 +1 2616 + 0058 - 4095 - 0130 - 0157 + 0139 + 0165 + 0154 + 0431 + 0058 - 6850 - 9232 29	-1. 9370 -2. 1732 +. 0057 6429 0233 0293 +. 0111 0055 +. 0117 +. 0154 +. 0066 3943 +3. 2070 -1. 9480	-1. 9370 -2. 0730 +. 0069 6256 0142 0260 +. 0153 +. 0016 +. 0206 +. 0180 +. 0074 3641 +3. 5431 -1. 4276 67
		·	$+1.2773$ $x_{13}=$	5445 +. 42629	-, 2115 +, 16558	+1. 13067	+. 72254	-1. 9480 +1. 52509	+1. 11697

	····	1					i i
<u> </u>	24	25	26	η (Lat.)	Σ (Lat.)	η (Long.)	Σ (Long.)
22 23 11 13	+4. 15 9835 5036 0178 2321	-0. 6223 0841 0156 0902	-1. 25	0 +0.3192 9112 +.1840 6156	0 +0.3643 8726 +.2289 3934	0 +0.4175 5024 +.4287 8304	0 +0.4626 4638 +.4736 6082
"	+2.4130	8122 +. 33659	-1. 25 +. 51803	-1.0236 +.42420	6728 +. 27882	4866 +. 20166	1358 +. 05628
	x24=	+3. 35 3029	-1.45	+2.7495	+2.7495	-1.6630	-1.6630
	21 22	0038 3937		+. 1161 +. 0831 +. 2020	+. 2750 +. 0884 +. 2305	+. 2094 +. 0317 +. 2642	+. 3684 +. 0370 +. 2927
	23 11	0141 0136		1522 +. 1609	1458 +. 2002	0839 +. 3748	0775 +. 4141
	13 24	0350 2734	42 07	2391 3445	1528 2265	3225 1638	2362 0457
		+2.3135 x ₂₅ =	-1.8707 +.80860	+2.5758 -1.11338	+3.0185 6 -1.30478	+1.3531 +.58487	9102 3 +. 39347
		24 25	+2.70 6475 -1.5126	-1. 9575 5303 +2. 0828	-1. 9575 3485	+1.3630 2521 -1.0941	+1. 3630 0703
		-	+. 5399	4050	+2.4408 +.1348 9	+.0168	7361 +. 556£ 7
1		ļ	x25=	+. 75014 +43. 4557	24986 +43. 4557	03112 +72. 6321	-1.03112 +72.6321
.			1 2 3	2359 0369 -1. 8536	+. 0295 0911 -1. 6801	- 2106 -2.1809 -5.6301	4614 -2, 5982 -5, 9324
			1 4	-4. 9061 -, 7866	-4. 7853 9098	-2.6859 0007	-2.7753 +.0028
			5 6 7	0274 0977	+. 0059 +. 0001	4356 0274	3027 +. 0244
			8 9	0135 -5. 7502	+. 0720 -6. 2462	0038 -20. 1312	+. 0415 -19. 2030
			10 12	2731 2092 -5. 0460	3411 2007	4059 4747 -7. 2954	4889 4619
			14 15 16	3566 -1. 0503	-6. 2441 2362 6174	3963 -1. 1644	-8.7360 2693 7086
			17 18	-3. 4149 0243	-2. 8206 0352	-8. 1300 0323	-7. 2129 0448
			19 20	0600 0445	1062 1054	0213 1448	+.0062 2547
			21 22	-1.8333 1036	-1. 9498 1182	2661 1773	-, 3105 -, 1964
			23 11	-1. 6485 -1. 8975 -1. 6329	-1.5787 -2.3606	5011 -10. 2945	4626 -11. 3731 -2. 1758
			13 24 25	4342 -2. 8678	-1. 0435 2854 -3. 3608	-2.9709 0981 7914	0274 5324
		_	25 26	3038	+. 1012	0005	0173
	ļ			+8. 5473	+8.5480 +46.3557	+8. 1609	+8. 1614 +75. 5321
]		2 3	0037 2252 -1, 5229		-1, 0106 -3, 0954 -6, 2509
				4 5	-4.6675 -1.0523		-2.8676 0126
				6 7	0013 0		2103 0217
				8 9	3828 -6. 7851		4535 -18. 3176
				10 12 14	- 4261 - 1926 -7, 7265		5888 4495 - 10. 4609
				15 16	1564 3629		1830 4312
	,			17 18	-2.3297 0509		-6.3993 0621
				19 20	1880 2497		0018 4481
				21 22	-2. 0738 1349 -1. 5118		3623 2176
				23 11 13	-2.9367		4270 -12. 5647 -1. 5936
				24 25	6668 1876 -3. 9386		0076 3582
				26	 0337		5740 +8. 1622
					+8. 5482		T-0, 1022

U. S. COAST AND GEODETIC SURVEY

Back solution for latitude

. 26	25	24	13	11		23	3	22	21	20
+0.75014 +.75014 +.075 +."074	-1. 11338 + . 60656 50682 051 "050	+0. 42420 +. 38860 17059 +. 64221 +. 064 +. "063	+1, 13067 - 08392 + 27377 +1, 32052 +, 132 +, "131	-0. 86847 -0.03733 +0.5409 +. 83217 -0.1954 002 "002		+0.86 00 +.22 +.44 00 +1.50 +.15 +."	3751 8454 5929 0198 0594	-0. 1596 1576 +. 3158 +. 0103 0006 +. 1782 +. 1860 +. 019 +. "019	69 01656 + . 02102 33 00192 55 + . 36237 21 + . 10541 99 25027 025	-0. 16579 - 21931 + .0320000261 + .026630412637034037"037
19	18	17	16	15		14		12	10	9
-0, 14945 +, 03064 -, 00267 +, 02843 -, 04732 -, 19158 -, 33195 -, 033 -, "033	-0.07686 +.04006 00380 +.04251 07594 00120 13484 21007 021 "021	+1. 48546 +. 01092 00058 +. 00379 19144 05562 00350 +1. 24903 +. 125 +. "124	+0.64002 +.01777 00094 +.00611 09063 00571 +.46264 +1.02926 +.103 +."102	+0. 52134 +. 80908 +. 18127 +1. 51169 +. 151 +. "149		-1. 49 + 36 + 43 - 69 - 00 "	5596 3671 9489 59	+0. 2131 +. 5741 0011 +. 6538 0121 +. 0053 +1. 4330 +. 143 +. "141	1 + .00516 600762 308013 8 + .03516 4 + .19133 508432 008	-1. 29049 +. 34421 01132 +. 02677 00192 93275 093 "092
8	7	6	5		4			3	2	1
+0.06543 +.26039 01867 10475 +.20240 +.020 +."020	+0. 20686 - 33121 + 08861 - 03574 - 004 - "004	+0. 1143 2225 0222 1304 013 "013	6 81 -0.08	900 900	+1.	74748 00107 00853 70563 03225 103 "'102	-	0. 73751 00173 01370 39654 +. 30296 +. 62844 +. 063 +. "062	-0.11016 00406 03221 +.30002 +.15359 +.015 +."015	+. 26537 00747 +. 07565 +. 33355 +. 033 +. 2"033

Back solution for longitude—Continued

26	25	24	13	11	23		22	21	20
-0.03112 03112 003 "004	+0.58487 02516 +.55971 +.056 +."084	+. 20166 01612 +. 18839 +. 37393 +. 037 +. "052	+1. 52509 +. 09268 +. 15940 +1. 77717 +. 178 +. "214	-2. 0228 +. 0412 +. 0315 +1. 1199 8302 083 "10	+ . 04 + . 16 + . 61 + . 61 + . 61 + . 11 + . 11	1143 3568 1812 3397 3320 —	-0. 20877 +. 17414 +. 18388 +. 01390 04021 +. 14002 +. 26296 +. 026 +. "037	-0. 27453 +. 01829 +. 02829 08178 +. 28471 +. 14896 +. 12394 +. 012 +. "016	-0. 29919 +. 24219 +. 04306 11075 +. 02092 +. 02043 08334 008 "012
19	18	17	16	15	14		12	10	9
+0.08895 +.04123 11329 +.02234 +.02234 04311 +.01955 +.002 +."003	-0. 08853 +. 05392 16129 +. 03340 +. 03761 00027 +. 00794 11722 012 "016	+2. 29200 +. 01470 02452 +. 00298 04308 +. 00328 00195 +2. 24341 +. 224 +. "335	+0.67391 +.02392 03992 +.00480 +.00534 00319 +.83096 +1.49582 +.150 +."207	+0. 5498 +1. 4533 +. 2634 +2. 2661 +. 227 +. "34	21 +. 55 14 +. 64 19 66 00	3185 5468 1414 61 -083	+0. 32100 +. 77264 04922 +. 51371 00679 +. 00776 +1. 55910 +. 156 +. "195	-0. 27825 +. 00695 32388 04471 +. 05110 +. 20816 38063 038 "049	+2 41462 + 46324 - 48089 + 03891 - 00866 +2 42722 + 243 + "287
8	7	6	5		4	3		2	1
+0. 03466 +. 37843 08426 +. 27258 +. 60141 +. 060 +. "'076	+0. 10962 +. 86188 +. 26329 +1. 23479 +. 123 +. "148	+0. 4556 +. 5791 +. 7684 +1. 8034 +. 180 +. "20	16 +.04 10 +1.46 19 +1.57 +.16	9610 5247 7874 58	-1. 29298 + 03712 + 11788 +1. 36020 + 22222 + 022 + "024	+++	28535 05964 18946 76439 06521 20665 021	-0. 84736 +. 14017 +. 44534 09865 36050 036 "042	-0. 25075 +. 25802 17756 17029 017 "021

U. S. COAST AND GEODETIC SURVEY

Latitude and longitude v's and total closures

Num- ber of		Latitude		p	. 1	Longitude		Total closure	Closure as
sec- tion	v	pv	pv 1		v	pv	102	in feet	part. One part in—
1	+0.33	+0.33	0. 11	1. 00	-0.17	-0.17	0. 03	3. 7	357, 000
2	18	30	. 05	1. 65	19	31	. 06	2. 6	292, 000
3	+.48	+.70	. 34	1. 45	+.15	+.22	. 03	5. 0	185, 000
4	+.40	+.40	. 16	1. 00	+.43	+.43	. 18	5. 9	224, 000
5	+.40	+.40	. 18	. 90	+.48	+.43	. 21	6. 6	• 220, 000
6	+. 17	+. 28	. 05	1. 65	12	- 20	. 02	2.1	390, 000
7	+. 69	+. 69	. 48	1. 00	+. 22	+ 22	. 05	7.2	183, 000
8	-1. 31	98	1, 28	. 75	72	- 54	. 39	14.9	120, 000
9	+. 90	+. 63	. 57	. 70	+. 20	+ 14	. 03	9.2	201, 000
10	+. 20	+. 22	. 04	1. 10	+. 60	+ 66	. 40	6.3	187, 00 0
11	+. 09	+. 10	. 01	1. 10	57	63	. 36	5. 8	205, 000
12	+. 24	+. 24	. 06	1. 00	63	63	. 40	6. 7	197, 000
13	80	40	. 32	. 50	+. 63	+. 32	. 20	10. 2	259, 000
14	+1. 00	+. 50	. 50	. 50	+. 31	+. 16	. 05	10. 4	254, 000
15	28	20	. 06	. 70	98	69	. 68	10. 2	186, 000
16	+. 06	+. 12	. 01	2.00	45	90	. 40	4.5	147, 000
17	+. 16	+. 32	. 05	2.00	+. 31	+. 62	. 19	3.5	189, 000
18	12	08	. 01	.70	45	32	. 14	4.6	413, 000
19	11	22	. 02	2.00	+. 22	+. 44	. 10	2.5	264, 000
20	26	23	. 06	.90	18	16	. 03	3.1	477, 000
21	69	55	. 38	. 80	61	49	. 30	9. 2	178, 000
22	35	23	. 08	. 65	03	02	. 00	3. 5	581, 000
23	42	34	. 14	. 80	59	48	. 28	7. 2	227, 000
24	+. 83	+. 66	. 55	. 80	+. 90	+. 72	. 65	12. 2	134, 000
25	+. 22	+. 18	. 04	. 80	+. 74	+. 59	. 44	7. 7	213, 000
26	26	22	. 06	. 85	03	03	.00	2.6	599, 000
27	13	26	. 03	2. 00	+. 26	+. 52	.14	2.9	227, 000
28	12	20	. 02	1. 65	+. 14	-+. 23	.03	1.8	440, 000
29	+. 22	+. 15	. 03	. 70	51	36	:18	5.5	344, 000
30	04	05	. 00	1. 25	10	12	.01	1.1	960, 000
31 32 33 34 35	06 04 +. 44 +. 45 87	05 05 +. 88 +. 90 78	.00 .00 :39 .40	. 80 1. 25 2. 00 2. 00 . 90	+. 73 +. 24 +. 14 +. 11 81	+. 58 +. 30 +. 28 +. 22 73	. 42 . 07 . 04 . 02 . 59	7.3 2.4 4.6 4.6 41.9	224, 000 440, 000 143, 000 143, 000 124, 000
36 37 38 39 40	+. 08 -1. 15 14 04 +. 57	+. 16 92 10 05 +. 46	.01 1.07 .01 .00 .26	2.00 .80 .70 1.20	38 +. 03 +. 64 +. 05 34	76 +. 02 +. 45 +. 06 27	. 29 . 00 . 29 . 00 . 09	3. 9 11. 5 6. 6 0. 6 6. 6	169, 000 142, 000 288, 000 2, 015, 000 248, 000
41	09	13	. 01	1. 45	+. 35	+. 51	. 18	3. 6	252, 000
42	+. 11	+. 14	. 02	1. 25	40	50	. 20	4. 1	258, 000
	Sun	1	8. 54				8. 17		

Changes in positions of junction points caused by the readjustment

Num- ber of junc- tion	Station		itud	ude and le result- le the re- lent	,long	itud Iron	ie and eresult- iold ad- t	New minus old
1	Carson	35 97	, 16 57	24. 922 32. 416	。 35 97	16 57	24. 848 32. 760	,, +0.074 344
2	Kyle	32 98	49 19	18. 039 11. 612	32 98	49 19	18. 014 12. 209	+. 02 5 597
3	Mission	29 98	42 09	52. 843 52. 109	29 98	42 09	52. 761 52. 996	+. 082 887
4	Donna	26 98	09 02	40. 593 44. 471	26 98	09 02	40. 459 45. 414	+. 134 943
5	Peters	29 100	10 36	32. 792 02. 238	29 100	10 36	32. 904 03. 100	112 862
6	Ingle	31 103	35 35	48. 685 24. 355	31 103	35 35	49. 018 25. 284	333 929
7	Phillips	. 34 104	59 07	03. 363 59. 629	34 104	59 07	03385 60. 204	022 575
8:	Aroya	38 103	48 10	09. 721 55. 580	38 103	48 10	10. 006 55. 610	285 030
9	Whitetank	33 112	34 33	01. 652 27. 501	33 112	34 33	02. 053 28. 731	401 -1. 230
10	Mount Nebo	39 111	48 45	38. 316 56. 235	39 111	48 45	39. 113 56. 918	797 683
11	Black	35 115	55 02	51. 689 35. 556	Not	prev	iously ad	justed.
12	Mount Diablo	37 121	52 54	54. 387 .47. 107	37 121	52 54	55. 482 48. 355	-1.095 -1.248
13	Tepusquet	34 120	54 11	36. 522 08. 335	34 120	54 11	37. 432 09. 654	910 -1. 319
14	Farmer	97	42 40	21. 212 34. 023	43 97	42 40	21. 389 34. 412	177 389
15	States	49 97	00 07	01. 447 39. 343	49 97	00 07	01. 687 39. 738	240 395
16	Sundance	44 104	28 27	44. 202 03. 099	44 104	28 27	44. 696 02. 821	−. 494 +. 278
17	Norge	48 103	53 47	38. 110 21. 176	48 103	53 47	38. 866 20. 556	756 +. 620
18	Oxford	42 112	16 05	10. 842 49. 460	42 112	16 05	11. 766 49. 972	924 512
19	Mount Ellis	45 110	34 57	38. 926 18. 432	Not	prev	iously ad	justed.
20	Goldstone	, 48 110	52 29	55. 681 18. 684	Not	pre	viously ad	justed.
21	Dry	44 117	10 39	08. 183 39. 844	44 117	10 39	09. 309 40. 845	-1. 126 -1. 001
22	Alder	45 119	50 56	59. 419 21. 080	45 119	50 56	60. 583 22. 229	-1. 164 -1. 149
-23	Bally	40 122	36 38	10. 879 59. 178	40 122	36 38	11. 939 60. 370	-1.060 -1.192
24	Red	45 121	56 49	06. 140 11. 100	45 121	56 49	07. 249 12. 344	-1. 109 -1. 244
25	Oroville	48 119	53 20	44. 203 12. 879	Not	prev	iously ad	justed.
26	Birch Point	48 122	56 49	30. 854 12. 021	48 122	56 49	31. 910 13. 395	-1. 056 -1. 374

CLOSURES FROM THE ADJUSTMENT

The loop closures are shown by the equations on page 16, but a better understanding of the closures can be had by computing them for the several individual loops. The figures in the equations represent in many cases the closures in overlapping loops and hence are not as enlightening as the results for the simple loops. In Figure 5 these individual loop closures are shown; the first number above the line gives the closure in meters, the second number is the approximate length of the loop in kilometers, and the number below the line shows the approximate proportional part that the closure bears to the whole length of the loop. Only 2 of these closures are greater than 1 part in 200,000, the largest one being 1 part in 162,000. The mean of all of the closures is approximately 1 part in 450,000. The closure around the entire outer boundary is 10.1 meters in a distance of some 8,504 kilometers, or approximately 1 part in 842,000. The total length of the arcs in the adjustment is approximately 20,000 kilometers.

Some idea of what these closures mean may be obtained when we consider that they are the result of a combination of various types of measurements which have been made under conditions in the field that are oftentimes very unfavorable. The base measurements form one element of control, the angle measurements serve to carry the length through the arcs from base to base, and finally the astronomic measurements furnish the azimuth control of the various arcs. Some of this work was done as much as 50 years ago, but the greater

part of it has been done in the past 25 years.

In the network included in this adjustment the length is controlled by 50 bases, and the orientation is maintained by 74 Laplace azimuths, both classes of control being more or less evenly distributed throughout the various sections of the net. The 50 bases have been measured within the past 50 years with various types of The older bases were measured with base measuring apparatus. bars or rods. In 1900-01 the measurement of nine bases along the ninety-eighth meridian was made with both bars and steel tapes, and in 1907 six bases were measured with both steel and invar tapes. Since 1907 all first-order bases have been measured with invar tapes. With these three types of apparatus different methods of measurement were employed. In spite of these facts the resulting loop closures demonstrate clearly that the general accord of bases is even much better than could be expected, and the fitting together of the various sections is satisfactory in every way.

In the previous adjustment of the transcontinental arc along the thirty-ninth parallel and of the arcs extending southward from this parallel, both along the ninety-eighth meridian as far as Texas and near the coast of California, the equations to hold the Laplace azimuths were not included. Because of this fact, the azimuths of the arcs became progressively bad and introduced errors in position that increased more and more as the distance from the starting point increased. On account of this swerving of the triangulation, as it is technically called, it was found very unsatisfactory to adjust the new arcs to close the loops, since all of the error of closure had to be absorbed by the new work. The introduction of Laplace azi-

muths to control the orientations of the arcs was an important step forward in geodetic work.

CHANGES IN POSITIONS

After the loop closures were determined and the equations given in the table on page 16 were formed, these equations were solved by the method of least squares to determine the best positions to be held in the junction figures. This solution resulted in the determination of the final geodetic position of a station in each one of the 26 junction figures. A list of these final positions is given on page 29. The positions of all of these same stations that had been previously adjusted are also given, together with the amount of the change of each brought about by the readjustment. These same changes are shown in Figure 7. It will be noted that junction No. 26 (see figs. 4 and 7) has changed -1.056 in latitude and -1.374 in longitude, or something like 30 meters in both latitude and longitude. This is the point that controls the data that have since been carried northward into western Canada and Alaska.

SECTION CLOSURES

Figure 6 and the table on page 28 give the amounts of latitude and longitude discrepancies that had to be absorbed by the various sections. They also give the total closure in length that had to be adjusted into each section and finally the approximate proportional part of the length of each section that this closure represents. It will be seen that no closure in position in any section is greater than 1 part in 120,000, and that the great majority of the sections have closures of 1 part in 200,000 or better. The mean of all of the closures of sections is approximately 1 part in 317,000. It is clear that these closures could be absorbed by the sections by the introduction of the latitude and longitude equations without any violence to the work in any of its parts. The whole network has therefore been fitted together in a rigid system without undue strain in any of its parts. Any short arc that may be observed in the future between sections of this framework should fit into the general scheme with comparatively small closure in position.

KIND OF FIGURES

'Special attention should be called to the fact that throughout the 20,000 kilometers of arcs only complete quadrilaterals or central point figures were used except in a single case in which at the last minute it was found necessary to reject one diagonal of a quadrilateral because of bad closures in the figure. It is believed that the excellence of the results developed in the closures of loops is due in large measure to the use of such figures in carrying the arcs over such great extent of territory. It is not necessary to do more than cite the results as found to justify such an opinion. It is the policy of this bureau to observe all first-order triangulation in this way whenever it extends such control into new territory. Experience certainly justifies such precautions in geodetic work that is to serve for basic control.

CONCLUSION

The loop closures resulting from this adjustment clearly show that the methods employed in the geodetic operations in this region are fully adequate for all purposes whether practical or scientific. The instructions under which these observations were made called for an average triangle closure of about 1 second with a maximum closure of 3 seconds except in a few cases; for Laplace azimuths to be observed every 6 to 10 figures in the arcs; and for bases to be measured when the sum of the R₁'s was somewhere between 80 and 100. These ideal conditions were not always rigidly observed, but the exceptions are few and in all cases due to unusual circumstances. Taken all in all, the loop closures obtained speak very highly of the excellence of the field work during the fifty-odd years in which the work was in progress. These results fully justify our belief that experience has taught this bureau a method of observation that meets all demands of accuracy for high-grade geodetic work.

The results of this adjustment show unquestionably that the method of observations is fully justified and also that the Bowie method of adjustment is economically and scientifically adequate for application to such complicated conditions. Much credit is due to the field officers of this bureau for the conscientious manner in which they have endeavored to attain the ideals of accuracy called for in their instructions. Much credit is also due to the office mathematicians for their efficient handling of the complicated

calculations that the adjustment demanded.

 $^{^1}$ The sum of the R_1 's is a function of the angles that carry the length through the chain of triangles See Special Publication No. 120, p. 5.